## Low-level laser treatment can reduce edema in second degree ankle sprains



Authors: Stergioulas A.

Source: J Clin Laser Med Surg., April 2004

## **Abstract**

Low-level laser therapy (LLLT) has been used for the last few years to treat sports injuries. The purpose of this study was to compare three therapeutic protocols in treating edema in second degree ankle sprains that did not require immobilization with a splint, under placebo-controlled conditions.

Forty-seven soccer players with second degree ankle sprains, selected at random, were divided into the following groups: The first group (n=16) was treated with the conventional initial treatment (RICE, rest, ice, compression, elevation), the second group (n=16) was treated with the RICE method plus placebo laser, and the third group (n=15) was treated with the RICE method plus an 820-nm GaA1As diode laser with a radiant power output of 40 mW at 16 Hz. Before the treatment, and 24, 48, and 72 h later, the volume of the edema was measured.

A three by three repeated measures ANOVA with a follow up post hoc test revealed that the group treated with the RICE and an 820-nm GaA1As diode laser presented a statistically significant reduction in the volume of the edema after 24 h (40.3 +/- 2.4 mL, p < 0.01), 48 h (56.4 +/- 3.1 mL, p < 0.002), and 72 h (65.1 +/- 4.4 mL, p < 0.001).

## Conclusion

LLLT combined with RICE can reduce edema in second-degree ankle sprains.

To view a list of all the clinical and financial benefits provided by Theralase® laser technology, please visit: www.theralase.com/benefits

To access abstracts and clinical studies related to chiropractic conditions, please visit: www.theralase.com/chiroclinical Request an in-clinic demonstration today and learn about the benefits that Theralase® laser therapy can bring to your practice.

Be part of the thousands of customers worldwide who use Theralase® therapeutic lasers to treat their patients.